

XIII. *Description of Brotera persica and Mustelia eupatoria, two new Plants cultivated in the Botanic Garden of Halle, by Curt Sprengel, M.D. Professor of Botany in the University of Halle.*

*Read March 3, 1801.*

BROTERA PERSICA.

TAB. XII.

CLASSI XIV. adnumeranda.

*Charaeter essentialis genericus.*

**L**ABII inferioris lacinia media cucullata, involvens genitalia, eaque cum impetu protrudens.

DESCRIPTIO.—*Herba quadripedalis, caule ramisque tetragonis, axillis (fig. a.) barbatis.*

*Folia* opposita, subsecunda, petiolata, ovata, ferrata, scabriuscula.

*Infloresc.* Cymata axillaris pedunculata. (fig. b. b.)

*Cal.* quinquearistatus, pubescens. (fig. c. c.)

*Corolla* minuta, calyce vix longior, ringens (fig. d.): labium superius bipartitum: (1.) inferius tripartitum, lacinia media (3.) cucullata, antheras cum stylo occultans, donec externus stimulus accedat, qui ilico deprimit cucullum, ut antheræ erectæ  
prodeant

prodeant. Corolla pallida, lutescit ad faucem cuculli: ipse cucullus rufus est.

*Stamina* pilosa. *Antheræ* luteæ. (fig. e.)

*Stylus* violaceus. *Stigma* capitatum. (fig. f.)

*Semina* quatuor oblonga nuda in fundo calycis. (fig. g.)

In honorem summi botanici Felicis Avellar Brotero, Prof. Conimbricensis, id genus dixi. Accepi autem femina ab amico *Thouino* e museo naturali Parisino, quo apportata sunt ab *Oliverio* et *Brugucro* e Persia.

#### MUSTELIA EUPATORIA.

#### TAB. XIII.

#### CLASSI XIX. adnumeranda.

#### *Charaeter essentialis genericus.*

*Anthodium* simplex polyphyllum. *Rec.* nudum. *Papp.* duplex, paleaceus & quinquearistatus. *Corollulæ* quinquefidæ.

DESCRIPTIO.—*Herba* bipedalis: caule inclinato, recurvato.

*Folia* opposita, fasciculata, oblonga, apice denticulata, lucida, punctata, trinervia.

*Inflor.* in corymbis, quorum medius primo floret, lateralibus elongatis ferius florescentibus.

*Anthodium* constanter quinqueflorum; polyphyllum, æquale. (fig. a.)

